

REFERENCES

- Borghesi A, Stronati M, Castagnoli R et al. Novel Approaches to the Study of Neonatal Infections. *Am J Perinatol.* 2018;35(6):570-574.
- Levy O. Innate immunity of the newborn: Basic mechanisms and clinical correlates. *Nat Rev Immunol.* 2007;7(05):379-390.
- Perez GF, Pancham K, Huseni S, Jain A, Rodriguez-Martinez CE, Preciado D et al. Rhinovirus-induced airway cytokines and respiratory morbidity in severely premature children. *Pediatr Allergy Immunol.* 2015;26:145–52.
- Darlow BA, Voss L, Lennon DR et al. Early-onset neonatal group B streptococcus sepsis following national risk-based prevention guidelines. *Aust N Z J Obstet Gynaecol.* 2016;56:69-74.
- Singh T, Barnes EH, Isaacs D; Australian Study Group for Neonatal Infections. Early-onset neonatal infections in Australia and New Zealand, 2002-2012. *Arch Dis Child Fetal Neonatal Ed.* 2019; 104(3):F248-F252.
- Darmstadt GL, Marchant T, Claeson M et al. A strategy for reducing maternal and newborn deaths by 2015 and beyond. *BMC Pregnancy Childbirth.* 2013;13:216.
- World Health Organization. World Health Report 2005: Make Every Mother and Child Count. Geneva: World Health Organization; 2005.
- Yoon HS, Shin YJ, Ki M. Risk factors for neonatal infections in full-term babies in South Korea. *Yonsei Med J.* 2008;49(4):530-536.
- UNICEF. The neonatal period is the most vulnerable time for a child. UNICEF, Neonatal Mortality, 2016. [https:// data. unicef. org/ topic/ child- survival/ neonatal- mortality/](https://data.unicef.org/topic/child-survival/neonatal-mortality/) (accessed 21 Aug 2017).
- Fleischmann-Struzek C, Goldfarb DM, Schlattmann P, Schlapbach LJ, Reinhart K, Kissoon N. The global burden of paediatric and neonatal sepsis: A systematic review. *Lancet Respir Med.* 2018; 6(3):223-230.
- Cnattingius S, Norman M, Granath F, Petersson G, Stephansson O, Frisell T. Apgar Score Components at 5 Minutes: Risks and Prediction of Neonatal Mortality. *Paediatr Perinat Epidemiol.* 2017; 31(4):328-337.
- Simonsen KA, Anderson-Berry AL, Delair SF, Davies HD. Early-onset neonatal sepsis. *Clin Microbiol Rev.* 2014;27(1):21-47.
- Lim JW, Chung SH, Kang DR, Kim CR. Risk Factors for Cause-specific Mortality of Very-Low-Birth-Weight Infants in the Korean Neonatal Network. *J Korean Med Sci.* 2015.
- Haque K. Definitions of bloodstream infection in the newborn. *Pediatr Crit Care Med.* 2005;6(3 Suppl):S45-9.
- National Institute for Health and Care Excellence (NICE). Antibiotics for neonatal infection. NICE quality standard 75. 2014; guidance. nice.org.uk/qs75, p.5.
- National Collaborating Centre for Women's and Children's Health. Antibiotics for early-onset neonatal infection: antibiotics for the prevention and treatment of early-onset neonatal infection. *Royal College of Obstetricians and Gynaecologists.* 2012.
- Maródi L. Neonatal innate immunity to infectious agents. *Infect Immun.* 2006 Apr;74(4):1999-2006.
- Velaphi S, Siegel JD, Wendel GD Jr, Cushion N, Eid WM, Sanchez PJ. Early-onset group B streptococcal infection after a combined maternal and neonatal group B streptococcal chemoprophylaxis strategy. *Pediatrics.* 2003;111(3):541-7.
- NICE clinical guideline. Antibiotics for early-onset neonatal infection: Antibiotics for the prevention and treatment of earlyonset neonatal infection. Available from: [http://www.nice.org.uk/ guidance/CG149](http://www.nice.org.uk/guidance/CG149), date of publication: August 2012.
- Escobar GJ, Puopolo KM, Wi S et al. Stratification of risk of early-onset sepsis in newborns ≥ 34 weeks' gestation. *Pediatrics.* 2014;133(1):30-36.
- Brown JVE, Meader N, Wright K, Cleminson J, McGuire W. Assessment of C-Reactive Protein Diagnostic Test Accuracy for Late-Onset Infection in Newborn Infants: A Systematic Review and Meta-analysis [published online ahead of print, 2020 Feb 3]. *JAMA Pediatr.* 2020;174(3):260-268.
- Ahmed E, Rehman A, Ali MA. Validation of serum C-reactive protein for the diagnosis and monitoring of antibiotic therapy in neonatal sepsis. *Pak J Med Sci.* 2017;33(6):1434-1437.
- Gutbir Y, Wainstock T, Sheiner E et al. Low Apgar score in term newborns and long-term infectious morbidity: A population-based cohort study with up to 18 years of follow-up. *Eur J Pediatr.* 2020; 179(6):959-971.
- Razaz N, Cnattingius S, Joseph KS. Association between Apgar scores of 7 to 9 and neonatal mortality and morbidity: Population based cohort study of term infants in Sweden. *BMJ* 2019;365:11656.

VOVI

None declared

financial support: none declared