

Phyto-activity of *Syzygium aromaticum* extract against pathogenic bacteria isolated from Chronic Tonsillitis Patients

By Hayder Fadhil Okab

Phyto-activity of *Syzygium aromaticum* extract against pathogenic bacteria isolated from Chronic Tonsillitis Patients

Iqbal Aziz Ameen ^{1*}, Hayder Fadhil Okab²

¹Department of Biology, Collage of Education for Girls, Al-Shatrah University, Thi-Qar, Iraq

²Collage of Education for Girls, Al-Shatrah University, Thi-Qar Health Directorate, Iraq

Corresponding author

m31947683@gmail.com

al-kalibyiqbal@utq.edu.iq

Abstract

Because of its antibacterial and antioxidant characteristics, the naturally occurring spice *Syzygium aromaticum* (Clove) is used as a medicine and a preservative. Although it is said to have originated in Indonesia, clove is now grown in numerous coastal regions at higher altitudes all over the world. Clove has been used for ages for its medicinal properties, which have been shown effective against a wide variety of illnesses. Clove has been found to be effective as an antibiotic, antiviral, anti-inflammatory, hepatoprotective, stress-reducing, interoceptive, and anesthetic. Clove's larvicidal efficacy against dengue is a novel use. Extraction of clove's volatile oil has found commercial usage in the creation of pharmaceutical, nutritional, and personal care products. Eugenol is a crucial phenolic molecule found in clove oil's various phytoconstituents. With regard to the issues raised, this study carried for detection activity¹⁸ of clove on pathogenic *S. aureus* and *S. pyogen* isolated from patients with chronic tonsillitis. The present study noted the activity of methanolic extract was higher activity than aqueous extract in all concentration with except 1% concentration was have not activity against both bacteria. Also, noted the *S. aureus* was has high susceptibility for both extract than *S. pyogen*.

Key words: *Syzygium aromaticum*, EO, Clove, *S. aureus*, *S. pyogen*

Introduction

As knowledge in the fields of nutrition and medicine expands, plants play an ever-increasing role in the lives of humans. Flavoring agents, culinary⁵ additives, coloring agents, preservatives, and medications are just few of the many uses for the dried root, seed, bark, fruit, or flowers of plants known as spices (Sachan *et al.*, 2018). The discovery of spices, which have been used for flavoring since prehistoric times, was a cause for celebration. ¹² Essential oils are described as an aromatic product of complex composition that is extracted from a botanically defined raw vegetable material using hydro-distillation, steam distillation¹⁴ or a suitable mechanical technique, as stated in the European Pharmacopoeia (Machadora *et al.*, 2020). Clove, or *Syzygium aromaticum*⁵¹, is one of the world's most expensive spices and the second most traded spice overall. *Caryophyllus aromaticus*, *Caryophyllus silvestris*, *Eugenia caryophyllus*, *Jambosa caryophyllus*, and *Myrtus caryophyllus* are all names for the clove tree (Kaur and Kaushal, 2019). Cloves, which are native to North Maluku Islands in Indonesia but are harvested at various⁵ times around the world, are widely cultivated, and can be found in stores year-round. Some wild clove variants are located in Bacan, Ternate, Motir, Tidore, Makian, and western sections of Irian Jaya, where⁸ the majority of cultivated clove comes from Indonesia, Madagascar, and Pemba. Only the states of Ka⁸ataka, Tamil Nadu, and Kerala are able to support clove growing in India. After Indonesia, India is now the world's largest user⁸ of clove. To put it simply, clove is a fragrant tree (Ginigaddara, 2018; Kaur and Kaushal, 2019). The word clove comes from the French words for nail (clou) and clove (clove). The clove tree, also known as the conical myrtle, reaches a maximum height of 10–12 meters and has a straight trunk (Khan, 2022). The dense, grayish, semi-erect branches are a dark shade of green. The large, opposing

5

leaves are oblong to elliptic in shape, simple obovate in outline, hairless, and covered in oil glands. Flowering on this tree starts at the 7-year mark and lasts for the next 80 years. Oils extracted from clove stems, clove buds, and clove leaves are all commercially accessible. The chemical make-up, aroma, and color of clove essential oils vary widely. The quantity of secondary metabolites present in clove essential oil is influenced by environmental, climatic, agricultural, and genetic variables (Nair, 2021).

Inflammation of the palatine tonsils, or tonsillitis, is most commonly (if not always) brought on by bacteria. Tonsillitis episodes are marked by high body temperature, a sore throat, odynophagia, swollen tonsils with or without exudate, and cervical lymphadenopathy (Klagisa *et al.*, 2022). Clinical diagnosis of tonsillitis relies on anamnestic history and physical examination. (Sydorchuk, 2018). When more than two bouts of tonsillitis are experienced within a 12-month period, the possibility of recurrent tonsillitis should be explored (Wilson *et al.*, 2023). Antibiotics are the standard treatment for tonsillitis episodes. An abscess in the peritonsillar space can develop if inflammation from the palatine tonsil spreads to the neighboring peritonsillar tissue (Klagisa *et al.*, 2022). The most typical purulent consequence of acute tonsillitis is a peritonsillar abscess (PTA). Clinical history and physical examination are used to confirm a PTA diagnosis. Microbiological analysis results show little variation between those with tonsillitis and healthy controls in studies comparing the two groups (Sanmark *et al.*, 2020). Without including comparisons to other patient groups, many studies just look at the microbiological outcomes of PTAs (Montero *et al.*, 2017; Klagisa *et al.*, 2022). The reasons for the failure of conservative therapeutic techniques in cases of tonsillitis are not well understood, and there are not many studies available on the topic. Research shows that *S. aureus* is highly prevalent in tonsillar samples from people who have tonsillitis (Smith and Heubi, 2018). The most common causes of tonsillitis are the bacteria *Staphylococcus aureus* and *Streptococcus pyogenes*. It is unknown, however, what part these organisms play in the development of abscesses, in the development of resistance to antibacterial therapy, and in the etiology of tonsillitis aggravation. It is important to investigate alternative defensive mechanisms, such as biofilm formation, because *S. aureus* does not exhibit great resistance to antibiotics in tonsillitis, in contrast to *S. pyogenes*. Antibiotic resistance has been linked to the development of biofilms (Katkowska *et al.*, 2020; Michalik *et al.*, 2020). Clove essential oil's antibacterial action in pasteurized milk has been documented against *S. aureus* and *L. monocytogenes*, among others (Radünz *et al.*, 2019). Clove oil was found to be highly effective against *Penicillium sp.*, *Aspergillus flavus*, and *S. aureus*, all of which were isolated from dried fish (*Decapterus maruadsi*) (Chouhan *et al.*, 2018). Clove oil's antibacterial properties were tested on *L. innocua*, *C. diversiformis*, and *S. aureus*, three gram-positive bacteria, and four gram-negative bacteria, via the broth microdilution method: *S. typhimurium*, *E. coli*, *S. liquefaciens*, and *S. putrefaciens* (de Meneses *et al.*, 2019).

Materials and Method

Plant Collection

Two fresh plant parts of *Syzygium aromaticum* was collected from thi-qar province markit, evaluated for their antimicrobial activity against two pathogenic bacteria were isolated from tonsillitis patients.

Phytochemical extraction Drying

The newly gathered plant components were cleaned in running water and then in sterile distilled water before being used in the extraction process. The substance was dried in a 50 degree Celsius oven for 48 hours before being ground into a powder (Köprü *et al.*, 2020).

Preparation of Bacterial Suspension

The tests bacteria *S. aureus* and *S. pyogen* were isolated from Nasiriyah city in Thi-Qar province. By adding sterile distilled water, the concentration of two harmful bacteria was brought up to the 0.5 McFarland standard (1.5×10^8 CFU/ml). Adjusting the turbidity of a microbiological suspension against McFarland standards ensures that the total number of microorganisms remains within an acceptable range. After thoroughly mixing 9.95 ml of 0.18M H₂SO₄ (1.0% w/v) with 0.05 ml of barium chloride (BaCl₂) (1.17% w/v BaCl₂.2H₂O), we had our 0.5 McFarland standard. The McFarland standard tube could be sealed snugly for up to 6 months to prevent evaporation loss. The test and the standard were shown on a white backdrop separated by a black line to facilitate comparison (Julianti *et al.*, 2017).

Preparation of Aqueous and Methanolic Extracts

The extraction of certain plant parts yields an extract, which is a combination of phytochemicals (Kazlauskaite *et al.*, 2022). The phytochemicals in the plant were extracted using a solvent, specifically 95% ethanol and distilled water. In order to create 100 ml of ethanol extract (1%, 5%, 10%, 15%, 20%, and 25% w/v), 1, 5, 10, 15, 20, and 25 g of powdered plant material were dissolved in adequate sterile ethanol and distilled water, respectively. For 24 hours, the mixture was left undisturbed in a sterile flask covered with aluminum foil to prevent evaporation, before being filtered using sterilized Whatman no.1 filter paper. The filtered extract was then evaporated in a water bath until only 1, 5, 10, 15, 20, and 25 milliliters of the original extract remained. Extracts prepared in this way, both ethanolic and aqueous, were tested right away for antibacterial activity using the agar well diffusion technique.

Statistical Analysis

The current data were statistically analysis by statistical software program SPSS version 26, based in using One way ANOVA for mean comparison, least significant difference LSD for mean difference, independent sample t test, and Scatter blot for draw the activity diagrams

Results

Antimicrobial Activity of Aqueous Extract on *S. aureus* and *S. pyogen* According to Concentrations

The present study showed the activity of aqueous extracts were increasing with concentration increasing against both *S. aureus* and *S. pyogen*. Also, noted the 1% concentration not-exhibit activity against *S. aureus*, while both 1% and 5% concentrations not-exhibit activity against *S. pyogen*, the study also, recorded the high activity against *S. aureus* was 13.3 ± 0.26 , while against *S. pyogen* 11.2 ± 0.23 . in the other hand the study noted the *S. aureus* bacteria effects more than *S. pyogen* in all concentrations with except 1% concentration at $p. value < 0.05$, as in table 3-1.

Table 3-1: Antimicrobial activity of aqueous extract on *S. aureus* and *S. pyogen* according to concentrations

Concentrations	<i>S. aureus</i>	<i>S. pyogen</i>	<i>p. value</i>
	Aqueous Extract Inhiation Zone Mean ± SD		
1%	0.00 ± 0.00 ^f	0.00 ± 0.00 ^e	1.000
5%	0.76 ± 0.30 ^e	0.00 ± 0.00 ^e	< 0.001
10%	3.66 ± 0.37 ^d	1.70 ± 0.36 ^d	< 0.001
15%	10.1 ± 0.58 ^c	4.86 ± 0.28 ^c	< 0.001
20%	11.9 ± 0.40 ^b	9.46 ± 0.20 ^b	< 0.001
25%	13.3 ± 0.26 ^a	11.2 ± 0.23 ^a	< 0.001
p. value	< 0.001	< 0.001	
LSD	0.65	0.40	

Antimicrobial Activity of Methanolic Extract on *S. aureus* and *S. pyogen* According to Concentrations

The present study showed the activity of aqueous extracts were increasing with concentration increasing against both *S. aureus* and *S. pyogen*. Also, noted the 1% concentration not-exhibit activity against *S. pyogen*, the study also, recorded the high activity against *S. aureus* was 15.5 ± 0.40 , while against *S. pyogen* 14.4 ± 0.75 . in the other hand the study noted the *S. aureus* bacteria effects significantly more than *S. pyogen* in 1%, 5%, 15% and 20% concentrations, while non-significantly in 10% and 25% concentrations at $p. value < 0.05$, as in table 3-2.

Table 3-2: Antimicrobial activity of methanolic extract on *S. aureus* and *S. pyogen* according to concentrations

Concentrations	<i>S. aureus</i>	<i>S. pyogen</i>	<i>p. value</i>
	Methanolic Extract Inhiation Zone Mean ± SD		
1%	1.06 ± 0.11 ^f	0.00 ± 0.00 ^f	1.000
5%	2.96 ± 0.32 ^e	2.40 ± 0.34 ^e	0.030
10%	5.96 ± 0.72 ^d	4.63 ± 0.60 ^d	0.072
15%	12.2 ± 0.58 ^c	7.70 ± 0.36 ^c	< 0.001

20%	14.1 ± 0.56 ^b	12.2 ± 0.25 ^b	0.006
5%	15.5 ± 0.40 ^a	14.4 ± 0.75 ^a	0.088
p. value	< 0.001	< 0.001	
LSD	0.88	0.81	

A Comparison between Aqueous and Methanolic Extracts on *S. aureus* According to Concentrations

The methanolic extract was exhibit a significant antimicrobial activity against *S. aureus* more than aqueous extract in all extract concentration, at p. value < 0.05, as in table 3-3.

Table 3-3: A Comparison between aqueous and methanolic extracts on *S. aureus* according to concentrations

Concentrations	Aqueous	Methanolic	p. value
	<i>S. aureus</i> Response Mean ± SD		
1%	0.00 ± 0.00	1.06 ± 0.11	< 0.001
5%	0.76 ± 0.30	2.96 ± 0.32	0.001
10%	3.66 ± 0.37	5.96 ± 0.72	0.008
15%	10.1 ± 0.58	12.2 ± 0.58	0.012
20%	11.9 ± 0.40	14.1 ± 0.56	0.005
25%	13.3 ± 0.26	15.5 ± 0.40	0.001

A Comparison between Aqueous and Methanolic Extracts on *S. pyogen* According to Concentrations

The methanolic extract was exhibit a significant antimicrobial activity against *S. pyogen* more than aqueous extract in all extract concentration, with except 1%, was not exhibit significant difference at p. value < 0.05, as in table 3-4.

Table 3-4: A Comparison between aqueous and methanolic extracts on *S. aureus* according to concentrations

Concentrations	Aqueous	Methanolic	<i>p. value</i>
	<i>S. pyogen</i> Response Mean ± SD		
1%	0.00 ± 0.00	0.00 ± 0.00	1.00
5%	0.00 ± 0.00	2.40 ± 0.34	< 0.001
10%	1.70 ± 0.36	4.63 ± 0.60	0.002
15%	4.86 ± 0.28	7.70 ± 0.36	0.001
20%	9.46 ± 0.20	12.2 ± 0.25	< 0.001
25%	11.2 ± 0.23	14.4 ± 0.75	0.002




Figure 3-1: the scatter blot of aqueous and methanolic extracts show the activity of extract increased with concentrations increasing, also exhibit the low and high activity against both bacteria.

Discussion

The current study noted the Phyto activity of clove methanolic extract was higher activity than clove aqueous extract in all concentration with except 1% concentration was have not activity against both bacteria. Also, recorded the *S. aureus* was has high susceptibility for both extract activity than *S. pyogen*.

The emergence of antibiotic-resistant bacteria in poor countries was triggered by both the infections themselves and the greater availability of some medications over the counter. Incorrect

dosing and the inappropriate use of antibiotics for viral infections may also play a role. Bacterial susceptibility ¹⁹ different antibiotics may exhibit geographical and temporal variations (Gebremariam et al., ¹⁵ 2015). Clove essential oil offers antimicrobial activity against a wide variety of infections. The -OH groups in the meta and ortho locations of the primary chemical composition have been linked to the antibacterial action. The cytoplasmic membrane of microorganisms can interact with these functional groups (Rajkowska et al., 2016; Behbahani et al., 20119). The study of Ayushi et al., (2020), was detected the clove oil extract has showed to ²² effective against *Staphylococcus SPP* and *A. niger*, also, noted the germicidal ¹⁷ activities against *S. aureus*, *P. aeruginosa*, and *K. pneumonia* have been shown by dispensing clove oil in 0.4% v/v concentrated with sugar solution. Study of Mohamed et al., (2018), involved 200 patients with corneal ulcer, their study showed 81.3% of activity of clove extract against pathogenic bacteria, also noted 97.5% of *S. aureus* inhibited by ³ clove extract, and 76% of *S. aromaticum*. The study of Ginting et al., (2021), involved extracts of clove and ³ cinnamon against gram negative bacteria isolated from chicken meat and their results revealed the effect of clove and cinnamon essential oils on both *E. coli*, *K. pneumoniae*, and ESBL-producing *E. coli* from ²¹ chicken meat observed through SEM showed changes ³ in bacterial cell structure. in studies of Abdel-Moaty et al., (2016), and Qian et al., (2020), we tested the antibacterial activity of clove and cinnamon essential oils against ESBL-producing bacteria as a comparison to the previous studies, and reported the essential oil of both plants have higher activity than antibiotics.

There may be a number of factors contributing to the recent uptick in research and usage of medicinal plants as alternative medicines in the prevention of infections. To begin, secondary metabolites from plants have enormous promise as antibacterial agents. Second, essential oils (EOs) can kill microorganisms without the harmful side effects of synthetic antibiotics. One more benefit of EOs is their ability to kill Gram-positive bacteria across the board, resulted from ⁷ the research. This detrimental effect on cellular permeability has been repeatedly described as the mechanism of action of *S. aromaticum* oil in studies of *S. aureus* and *S. pyogen* in recent years (Xu et al., 2016), which agrees with the negative impact of *S. aromaticum* oil on *S. aureus* cellular permeability found here.

References

- Abdel-Moaty, M. M., El-Hendawy, H. H., Mohamed, W. S., & Abdel-All, S. M. (2016). A potential role of cinnamon bark essential oil and its major component in enhancing antibiotics effect against clinical isolates of extended-spectrum beta-lactamase producing *Escherichia coli*. *Der Pharm. Lett*, 8(7), 138-147.
- Ayushi, U. A., Danish, S. M., & Mohammad, U. P. (2020). A review on biological and therapeutic uses of *Syzygium aromaticum* Linn.(Clove): Based on phyto-chemistry and pharmacological evidences. *International Journal of Botany Studies*, 5, 33-39.
- Behbahani, B. A., Noshad, M., & Falah, F. (2019). STUDY OF CHEMICAL STRUCTURE, ANTIMICROBIAL, CYTOTOXIC AND MECHANISM OF ACTION OF *SYZYGIUM*

AROMATICUM ESSENTIAL OIL ON FOODBORNE
PATHOGENS. *Potravinarstvo*, 13(1).

- Chouhan, S., Sharma, K., & Guleria, S. (2017). Antimicrobial activity of some essential oils—present status and future perspectives. *Medicines*, 4(3), 58.
- de Meneses, A. C., Sayer, C., Puton, B. M., Cansian, R. L., Araújo, P. H., & de Oliveira, D. (2019). Production of clove oil nanoemulsion with rapid and enhanced antimicrobial activity against gram-positive and gram-negative bacteria. *Journal of Food Process Engineering*, 42(6), e13209.
- Gebremariam, T. T., Alemu, T. A., & Daba, K. T. (2015). Bacteriology and risk factors of bacterial keratitis in Ethiopia. *Health Science Journal*, 9(5), 1.
- Ginigaddara, G. A. S. (2018). Wonders of spices in Sri Lanka. *Indian Spices: The Legacy, Production and Processing of India's Treasured Export*, 63-93.
- Ginting, E. V., Retnaningrum, E., & Widiasih, D. A. (2021). Antibacterial activity of clove (*Syzygium aromaticum*) and cinnamon (*Cinnamomum burmannii*) essential oil against extended-spectrum β -lactamase-producing bacteria. *Veterinary world*, 14(8), 2206.
- Julianti, E., Rajah, K. K., & Fidrianny, I. (2017). Antibacterial activity of ethanolic extract of cinnamon bark, honey, and their combination effects against acne-causing bacteria. *Scientia pharmaceutica*, 85(2), 19.
- Katkowska, M., Garbacz, K., Kopala, W., Schubert, J., & Bania, J. (2020). Genetic diversity and antimicrobial resistance of *Staphylococcus aureus* from recurrent tonsillitis in children. *Apmis*, 128(3), 211-219.
- Kaur, K., & Kaushal, S. (2019). Phytochemistry and pharmacological aspects of *Syzygium aromaticum*: A review. *Journal of Pharmacognosy and Phytochemistry*, 8(1), 398-406.
- Kazlauskaite, J. A., Ivanauskas, L., Marksa, M., & Bernatoniene, J. (2022). The Effect of Traditional and Cyclodextrin-Assisted Extraction Methods on *Trifolium pratense* L.(Red Clover) Extracts Antioxidant Potential. *Antioxidants*, 11(2), 435.
- Khan, S. (2022). Origin, taxonomy, and benefits of *Syzygium aromaticum*. *Authorea Preprints*.
- Klagisa, R., Racenis, K., Broks, R., Balode, A. O., Kise, L., & Kroica, J. (2022). Analysis of Microorganism Colonization, Biofilm Production, and Antibacterial Susceptibility in Recurrent Tonsillitis and Peritonsillar Abscess Patients. *International Journal of Molecular Sciences*, 23(18), 10273.
- Köprü, S., Say, R., Karaman, K., Yilmaz, M. M., & Kaplan, M. (2020). Optimization of processing parameters for the preparation of clove (*Syzygium aromaticum*) hydroalcoholic extract: A response surface methodology approach to characterize the

- biofunctional performance. *Journal of Applied Research on Medicinal and Aromatic Plants*, 16, 100236.
- Machadora, C. A. T., Lepiksonb, H. A., Antônio, M., de Andrade, N., Ferreirad, Y. T. S., & Santos, S. F. (2020). Essential Oil Extraction Industry: Exploring the Strategic Technology Dimension.
- Michalik, M., Samet, A., Podbielska-Kubera, A., Savini, V., Międzobrodzki, J., & Kosecka-Strojek, M. (2020). Coagulase-negative staphylococci (CoNS) as a significant etiological factor of laryngological infections: A review. *Annals of Clinical Microbiology and Antimicrobials*, 19(1), 1-10.
- Mohamed, M. S., Abdallah, A. A., Mahran, M. H., & Shalaby, A. M. (2018). Potential alternative treatment of ocular bacterial infections by oil derived from *Syzygium aromaticum* flower (Clove). *Current Eye Research*, 43(7), 873-881.
- Montero, E., Iniesta, M., Rodrigo, M., Marín, M. J., Figuero, E., Herrera, D., & Sanz, M. (2017). Clinical and microbiological effects of the adjunctive use of probiotics in the treatment of gingivitis: A randomized controlled clinical trial. *Journal of clinical periodontology*, 44(7), 708-716.
- Nair, K. P. (2021). Clove. *Minor Spices and Condiments: Global Economic Potential*, 127-142.
- Qian, W., Sun, Z., Wang, T., Yang, M., Liu, M., Zhang, J., & Li, Y. (2020). Antimicrobial activity of eugenol against carbapenem-resistant *Klebsiella pneumoniae* and its effect on biofilms. *Microbial pathogenesis*, 139, 103924.
- Radünz, M., da Trindade, M. L. M., Camargo, T. M., Radünz, A. L., Borges, C. D., Gandra, E. A., & Helbig, E. (2019). Antimicrobial and antioxidant activity of unencapsulated and encapsulated clove (*Syzygium aromaticum*, L.) essential oil. *Food chemistry*, 276, 180-186.
- Rajkowska, K., Nowak, A., Kunicka-Styczyńska, A., & Siadura, A. (2016). Biological effects of various chemically characterized essential oils: Investigation of the mode of action against *Candida albicans* and HeLa cells. *RSC advances*, 6(99), 97199-97207.
- Sachan, A. K., Kumar, S., Kumari, K., & Singh, D. (2018). Medicinal uses of spices used in our traditional culture: Worldwide. *Journal of Medicinal Plants Studies*, 6(3), 116-122.
- Sanmark, E., Wikstén, J., Välimaa, H., Aaltonen, L. M., Ilmarinen, T., & Blomgren, K. (2020). Peritonsillar abscess may not always be a complication of acute tonsillitis: A prospective cohort study. *PLoS One*, 15(4), e0228122.
- Smith, M. M., & Heubi, C. H. (2018). Infections of the Neck and Pharynx in Children. *Current Treatment Options in Pediatrics*, 4, 211-220.

Sydorchuk, A. S. (2018). PSEUDOMONAS AERUGINOSA AS CAUSATIVE AGENT OF ACUTE PYOGENIC TONSILLITIS IN YOUNG INDIAN FEMALE: CASE REPORT.

Wilson, J. A., O'Hara, J., Fouweather, T., Homer, T., Stocken, D. D., Vale, L., ... & Teare, M. D. (2023). Conservative management versus tonsillectomy in adults with recurrent acute tonsillitis in the UK (NATTINA): a multicentre, open-label, randomised controlled trial. *The Lancet*.

Xu, J. G., Liu, T., Hu, Q. P., & Cao, X. M. (2016). Chemical composition, antibacterial properties and mechanism of action of essential oil from clove buds against *Staphylococcus aureus*. *Molecules*, 21(9), 1194.

Phyto-activity of Syzygium aromaticum extract against pathogenic bacteria isolated from Chronic Tonsillitis Patients

ORIGINALITY REPORT

19%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.mdpi.com Internet	71 words — 3%
2	opensiuc.lib.siu.edu Internet	61 words — 2%
3	www.ncbi.nlm.nih.gov Internet	56 words — 2%
4	media.neliti.com Internet	44 words — 2%
5	www.phytojournal.com Internet	44 words — 2%
6	1library.org Internet	25 words — 1%
7	www.tandfonline.com Internet	25 words — 1%
8	"Indian Spices", Springer Science and Business Media LLC, 2018 Crossref	20 words — 1%
9	"Handbook of Nutraceuticals and Natural Products", Wiley, 2022	13 words — < 1%

10 docksci.com 13 words — < 1%
Internet

11 Tuqa A. Omeer, H. Mohammed, Basim Abdulhussein Jarullah, Safaa Halool Mohammed. 12 words — < 1%
"Enhancement the electronic and optical properties of the T-Graphene nanosheet in the present various impurities: DFT study", Chemical Physics Impact, 2023
Crossref

12 academic.oup.com 12 words — < 1%
Internet

13 digitalcommons.bau.edu.lb 12 words — < 1%
Internet

14 Muhammad Saeed, Muhammad S. Khan, Mahmoud Alagawany, Mayada R. Farag et al. 11 words — < 1%
"Clove (Syzygium aromaticum) and its phytochemicals in ruminant feed: an updated review", Rendiconti Lincei. Scienze Fisiche e Naturali, 2021
Crossref

15 José Nabor Haro-González, Gustavo Adolfo Castillo-Herrera, Moisés Martínez-Velázquez, Hugo Espinosa-Andrews. 10 words — < 1%
"Clove Essential Oil (Syzygium aromaticum L. Myrtaceae): Extraction, Chemical Composition, Food Applications, and Essential Bioactivity for Human Health", Molecules, 2021
Crossref

16 Niranjan Das, Akash Dey, Sudip Kumar Mandal, Debanjan Chatterjee, Rajan Logesh, Hari Prasad Devkota. 10 words — < 1%
"Secondary metabolites of clove (Syzygium aromaticum)", Elsevier BV, 2022
Crossref

17 Daljit S. Arora. "Some Indian Spices and Their Antimicrobial Properties", From Ethnomycology to Fungal Biotechnology, 1999
Crossref 9 words — < 1%

18 coek.info
Internet 9 words — < 1%

19 dspace.alquds.edu
Internet 9 words — < 1%

20 www.fag.esn.ac.lk
Internet 9 words — < 1%

21 jeb.co.in
Internet 8 words — < 1%

22 link.springer.com
Internet 8 words — < 1%

23 plantarchives.org
Internet 8 words — < 1%

24 Qing Liu, Xiao Meng, Ya Li, Cai-Ning Zhao, Guo-Yi Tang, Hua-Bin Li. "Antibacterial and Antifungal Activities of Spices", International Journal of Molecular Sciences, 2017
Crossref 6 words — < 1%

25 Renata Klagisa, Karlis Racenis, Renars Broks, Arta Olga Balode, Ligija Kise, Juta Kroica. "Analysis of Microorganism Colonization, Biofilm Production, and Antibacterial Susceptibility in Recurrent Tonsillitis and Peritonsillar Abscess Patients", International Journal of Molecular Sciences, 2022
Crossref 6 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES OFF
EXCLUDE MATCHES OFF